



Newly Diagnosed HIV Cases in the Philippines

In August 2014, there were 509 new HIV Ab sero-positive individuals confirmed by the STD/AIDS Cooperativge Central Laboratory (SACCL) and reported to the HIV and AIDS Registry (Table 1). This was 33% higher compared to the same period last year (n=382). [Figure 1]

Table 2. Percentage of HIV Cases per Region (August 2014)

Region	% of Cases
I	1%
II	1%
III	9%
IVA	19%
IVB	1%
V	1%
VI	5%
VII	10%
VIII	1%
IX	3%
X	2%
XI	6%
XII	3%
CAR	1%
CARAGA	1%
ARMM	<1%
NCR	37%

Most of the cases (96%) were male. The median age was 27 years (age range: 1-59 years). The 20-29 year (58%) age group had the most number of cases.

Reported modes of transmission were sexual contact (473), needle sharing among injecting drug users (34) and mother to child transmission (2) [Table 3, page 2]. Males having sex with other males (87%) was the predominant type of sexual transmission [Figure 2]. Most (92%) of the cases were still asymptomatic at the time of reporting [Figure 3].

In August 2014, 82% of the new HIV cases came from NCR, Region 4A, Region 7, Region 3 and Region 11 [Table 2].

Table 1. Quick Facts

Demographic Data	August 2014	Jan-Aug 2014	Cumulative 1984-2014
Total Reported Cases	509	3,908	20,424
Asymptomatic Cases	467	3,557	18,567
AIDS Cases	42	351	1,857
Males	491	3,758	18,418*
Females	18	150	1,995*
Youth 15-24yo	146	1,100	5,289
Children <15yo	2	3	68

*Note: No data available on sex for (11) cases.

Figure 1. Number of New HIV Cases per Month (2012-2014)




Figure 2. Comparison of the Proportion of Types of Sexual Transmission in 2014, 2013 & Cumulative Data (1984-2014)





Figure 3. Number of HIV/AIDS Cases Reported in the Philippines by Year, January 1984 to August 2014 (N=20,424)




Demographic Characteristics (1984-2014)

Ninety-six percent of the 3,908 cases in 2014 were male (3,758). Ages ranged from 1 to 82 years old (median 28 years). The 20-29 year age group had the most (58%) number of cases for 2014. For the male age group, the most number of cases were found among the 20-24 years old (25%), 25-29 years old (33%), and 30-34 years old (20%) [Figure 4].

From 1984 to 2014, there were 20,424 HIV Ab sero-positive cases reported (Table 1), of which 18,567 (91%) were asymptomatic and 1,857 (9%) were AIDS cases. As shown in Figure 4, there is a significant difference in the number of male and female cases reported. Ninety percent (18,418) were male. Ages ranged from 1-82 years (median 28 years). The age groups with the most number of cases were: 20-24 years (23%), 25-29 (31%), and 30-34 years (19%) [Figure 4].

Figure 4. Comparison of the Distribution of Male and Female HIV Cases by Age-Group and Certain Highlighted Years



*Note: 74 did not report age, 11 did not report sex, 10 did not report age and sex

Modes of Transmission (1984-2014)

In 2014, ninety-four percent (3,654) were infected through sexual contact, 6% (251) through needle sharing among injecting drug users and 3 was infected through mother to child transmission (Table 3). There were 3,514 males and 140 females infected through sexual transmission. The age range of those infected through sexual transmission was 16-82 years old (median 28 years).



Of the 20,424 HIV positive cases reported from 1984 to 2014, 93% (18,999) were infected through sexual contact, 5% (962) through needle sharing among injecting drug users, <1% (65) through mother-to-child transmission, <1% (20) through blood transfusion and needle prick injury <1% (3) [Table 3]. No data is available for 2% (375) of the cases.

Table 3. Reported Modes of HIV Transmission

Mode of Transmission	August 2014 n=509	Jan-Aug 2014 n=3,908	Cumulative N=20,424
Sexual Contact	473	3,654	18,999
<i>Heterosexual contact</i>	<i>60(13%)</i>	<i>546(15%)</i>	<i>4,620(24%)</i>
<i>Homosexual contact</i>	<i>243(51%)</i>	<i>1,926(53%)</i>	<i>8,804(46%)</i>
<i>Bisexual contact</i>	<i>170(36%)</i>	<i>1,182(32%)</i>	<i>5,575(29%)</i>
Blood/Blood Products	0	0	20
Injecting Drug Use	34	251	962
Needle Prick Injury	0	0	3
Mother-to-Child	2	3	65
No Data Available	0	0	375

Figure 6. HIV Transmission by Age-Group, 2014 (n=3,908)

Figure 5. Proportion of Types of Sexual Transmission, January 1984–August 2014




Geographic Distribution (1984-2014)

Since 1984 to present, there were 20,424 cases reported. Almost half (9,233) came from the National Capital Region. [Figure 7]. Thirteen percent (2,589) came from region 4A, followed by 10% (1,833) from Region 7, 8% (1,612) from Region 3, 6% (1,178) from Region 11 and the rest of the country comprises 15% (2,841) of all the cases.

**Note: 1,138 or 6% of the cases had no reported data on their address at the time of diagnosis*

Figure 7. Percentage of HIV Cases by Region, January 1984- August 2014



AIDS Cases (1984-2014)

Of the 3,908 HIV positive cases in 2014, three hundred fifty-one were reported as AIDS cases. Ninety seven percent (342) were male and 3% (9) were female. Ages ranged from 17-69 years (median 30 years). Three hundred forty-four acquired the infection through sexual contact (184 homosexual, 115 bisexual and 45 heterosexual) and 7 acquired through injecting drug use.

From 1984 to 2014, there were 1,857 AIDS cases reported. Eighty-four percent (1,563) were male and 16% (294) were female. Median age is 32 years (age range: 1-81 years). Sexual contact was the most common mode of HIV transmission, accounting for 96% (1,776) of all reported AIDS cases. Forty-four percent (784) of sexual transmission was through homosexual contact, followed by heterosexual contact (621) then bisexual contact (371).

Overseas Filipino Workers (1984-2014)


In 2014, there were 426 HIV positive OFWs, comprising 11% of cases reported for the year [Figure 8]. Of these, 376 (88%) were male and 50 (12%) were female. Most were infected through sexual contact (147 heterosexual, 170 homosexual and 108 bisexual).

There were 3,064 HIV positive OFWs since 1984, comprising 15% of all reported cases [Figure 8]. Eighty-two percent (2,498) were males. Ages ranged from 18 to 80 years (median 34 years). Sexual contact (98%) was the predominant mode of transmission (Table 4). Eighty-nine percent (2,714) were asymptomatic while 11% (350) were AIDS cases.

Table 4. Mode of HIV Transmission Among OFWs

Mode of Transmission	August 2014 n=47	Jan-Aug 2014 n=426	Cumulative N=3,064
Sexual Transmission	47	425	2,998
<i>Heterosexual contact</i>	<i>15(32%)</i>	<i>147(35%)</i>	<i>1,502(50%)</i>
<i>Homosexual contact</i>	<i>18(38%)</i>	<i>170(40%)</i>	<i>852(28%)</i>
<i>Bisexual contact</i>	<i>14(30%)</i>	<i>108(25%)</i>	<i>644(21%)</i>
Blood/Blood Products	0	0	10
Injecting Drug Use	0	1	5
Needle Prick Injury	0	0	3
No Data Available	0	0	48

Figure 8. Number of OFWs and the Proportion of OFWs among the Total HIV Cases by Year



Deaths among People with HIV

From January 1984 to August 2014, there were 1,052 reported deaths among people with HIV [Table 5]. Eighty-two percent (862) were male. In total, there has been 124 deaths among youth (15-24 years old) and 15 deaths among children (<15 years old).

The annual number of deaths from 1984-2010 ranges from 2 to 36 cases with a median of 14 deaths per year. Beginning 2011, there has been an increase in the number of recorded deaths. There were 68 in 2011, and 176 in 2012. The DOH-NEC established an official reporting form in 2012.

From January to August 2014, there were a total of 113 deaths. Ninety-six percent (109) were male [Table 5]. The highest number of deaths occurred in the 25-29 (41%) age group [Figure 9]. This was followed by the 30-34 (19%) and 20-24 (12%) and 35-39 (12%) age groups. For the month of August 2014, there were 12 deaths* [Table 5].

Among the reported deaths in 2014, sexual contact (95%) was the most common mode of HIV transmission (56 homosexual, 37 bisexual, 14 heterosexual). There were 6 cases who got infected through injecting drug use (5%) [Figure 9].


*This include deaths that fall under the reporting month only.

Table 5. Demographic data of reported deaths among People with HIV

Demographic Data	August 2014	Jan-Aug 2014	Cumulative* (1984-2014)
Total Reported Deaths	12	113	1052
Males	11	109	862
Females	1	4	190
Youth 15-24yo	0	15	124
Children <15yo	0	0	15

*Note: No year of death reported for 46 cases

Figure 9. HIV Transmission by Age-Group of Reported Deaths



PLHIV on Anti-Retroviral Therapy (ART)

As of August 2014, there were 7,380* People Living with HIV presently on Anti Retroviral Therapy. These are the combined numbers of adult and pediatric patients currently enrolled and accessing Anti-Retroviral drugs in the 19 treatment hubs listed on the right.

Treatment Hubs in the Philippines	
1.	Ilocos Training and Regional Medical Center
2.	Cagayan Valley Medical Center
3.	Baguio General Hospital and Medical Center
4.	Jose B. Lingad Medical Center
5.	James L. Gordon Memorial Hospital
6.	Makati Medical Center
7.	Philippine General Hospital
8.	Research Institute for Tropical Medicine
9.	San Lazaro Hospital
10.	The Medical City
11.	Bicol Regional Training and Teaching Hospital
12.	Corazon Locsin Montelibano Memorial Regional Hospital
13.	Western Visayas Medical Center
14.	Gov. Celestino Gallares Memorial Hospital
15.	Vicente Sotto Memorial Medical Center
16.	Zamboanga City Medical Center
17.	Southern Philippines Medical Center
18.	Northern Mindanao Medical Center
19.	Eastern Visayas Regional Medical Center

* This is not a cumulative number. It does not include those who already have died, left the country, lost to follow up or decided to stop taking

Blood Units Confirmed for HIV

As of August 2014, 285 blood units were confirmed positive for HIV by RITM. There is no available data yet on the total number of blood units donated.

These are confirmed positive blood units, not blood donors. One donor can donate more than one blood unit. HIV positive blood donors may not be in the HIV & AIDS Registry unless they underwent voluntary counseling and testing as individuals.

Table 6. Number of Confirmed HIV Positive Blood Units

Month	2014
January	40
February	29
March	45
April	34
May	27
June	25
July	40
August	45
September	
October	
November	
December	
Total	285



National HIV/AIDS & STI Strategic Information and Surveillance Unit

National Epidemiology Center, Department of Health, Bldg. 19, San Lazaro Compound, Sta. Cruz, Manila 1003 Philippines

Tel: +632 651-7800 local 2926, 2952
Fax: +632 495-0513
Email: HIVepicenter@gmail.com
Website: http://www.doh.gov.ph

Philippine HIV & AIDS Registry Report Editorial Team:

Patricia Lagayza
Patricia Lagayza T. Lagayza, RN, MSPH
HIV Surveillance Assistant, HIV Unit

Kristina Lopez
Kristina Lopez G. Bonifacio, BSW
Asst. HIV Surveillance Officer, HIV Unit

Noel S. Palapayan
Noel S. Palapayan, RN, NCM-ESP
HIV Surveillance Officer, HIV Unit

Enrique A. Torio
Enrique A. Torio, MD, PHSAE
Chief, SIME, NEC

Enrique A. Torio
Enrique A. Torio, MD, PHSAE, FRCO, CESO III
Director IV, NEC

Philippine HIV & AIDS Registry

The Philippine HIV & AIDS Registry is the official record of the total number of laboratory-confirmed HIV positive individuals, AIDS cases and deaths, and HIV positive blood units in the Philippines. All individuals in the registry are confirmed by the STD/AIDS Cooperative Central Laboratory (SACCL) at San Lazaro Hospital. While all blood units are confirmed by the Research Institute for Tropical Medicine (RITM). Both are National Reference Laboratories (NRL) of the Department of Health (DOH).

Mandatory HIV testing is unlawful in the Philippines (RA 8504). The process of reporting to the Registry is as follows: All blood samples from accredited HIV testing facilities that are screened HIV reactive are sent to SACCL (Individuals) or RITM (blood units) for confirmation. Confirmed HIV positive individuals and blood units are reported to the DOH-National Epidemiology Center (NEC), and are recorded in the Registry.

The Registry is a passive surveillance system. Except for HIV confirmation by the NRL, all other data submitted to the Registry are secondary and cannot be verified. An example would be an individual's reported place of residence. The Registry is unable to determine if this reported address is where the person got infected, or where the person lived after being infected, or where the person is presently living, or whether the address is valid. This limitation has major implications to data interpretation. Thus, readers are cautioned to carefully weigh the data and consider other sources of information prior to arriving at conclusions.