



Factsheet N°1: HIV Epidemic Update of Nepal, as of August 2010

Facts and Figures about HIV Epidemic in Nepal

- ☐ The first HIV infection was detected in 1988 in Nepal. Since then HIV and AIDS epidemic has evolved from low- to **concentrated** among **High Risk Groups**:
 - 🚫 Injecting drug users,
 - 🚫 Female sex workers,
 - 🚫 Clients of female sex workers,
 - 🚫 Men who have sex with men and
 - 🚫 Labour migrants.
- ☐ Heterosexual transmission is dominant.
- ☐ HIV prevalence in general population is <1%.

Situation of HIV in Nepal, 2009:

- ☐ Estimated number of HIV infections (2009) by age groups
 - 🚫 Children (0-14 years): 3,544
 - 🚫 Adults (15-49 years): 52,504
 - 🚫 Adults (50+ years): 7,480
 - 🚫 **Total: 63,528**
- ☐ Women living with HIV: 28.6%
- ☐ Cumulative number of reported HIV infections (1988 - 16 August 2010): 16,262


Source: NCASC, 2010

Table 1: Estimated HIV Infections by Risk Groups, 2009

Population sub-groups (15+ years)	Total infections	% share
↑ Injecting drug users	2,534	4.2
↑ Men who have sex with men	3,699	6.2
↑ Female sex workers	605	1.0
↑ Clients of sex workers	2,996	5.0
↑ Labour migrants	17,653	29.4
↑ Remaining low risk males	15,697	26.2
↑ Remaining low risk females	16,800	28.0
Total	59,984	100.0


Source: NCASC, 2010

Figure 1: Adult (15-49) HIV Prevalence, 2009



Source: NCASC, 2010

Figure 2: Number of Estimated HIV Infections vs. Estimated HIV Prevalence, 2009



Source: NCASC, 2010

For Further Information Please Contact:

National Centre for AIDS and STD Control, Teku, Kathmandu

Phone: +977 - 1 - 4261 653, 4362 753, 4258 219


Email: ncasc@mos.com.np

Fax: +977 - 1 - 4261 406

Website: www.ncasc.gov.np



Figure 3: Distribution of Estimated HIV Infections by Sub-Population Groups, 1980-2015



Source: NCASC, 2010

- ✘ Heterosexual transmission is the key route of spreading HIV in Nepal. About 90% infections are transmitted through sexual transmission.
- ✘ Injecting drug users, female sex workers and their clients and men who have sex with men are the key drivers of the epidemic.
- ✘ Further intensifying the effective targeted interventions for high risk groups with improved coverage is critical to contain the epidemic among high risk groups and to prevent spread into large general low risk population.

Table 2: Key Indicators of HIV and AIDS Situation in Nepal, 2009

Indicators	Value (2009)
✘ Adult (15-49) HIV prevalence	0.39% (0.3% - 0.5%)
✘ Proportion of women 15-49 living with HIV	28.6%
✘ Proportion of women and girls 15-24 living with HIV	6.2%
✘ Average number of new infections per year	4,760
✘ Average number of new infections per day	13
✘ Average number of new infections amongst children (0-14) per year	468
✘ Average number of AIDS deaths per year	4,701
✘ Average number of AIDS deaths among children (0-14) per year	292
✘ Average number of AIDS deaths among children (0-4) per year	106
✘ Total number of AIDS orphans till 2009	21,947
✘ Average number of AIDS orphans per year	455

Source: NCASC, 2010

For Further Information Please Contact:

National Centre for AIDS and STD Control, Teku, Kathmandu

Phone: +977 - 1 - 4261 653, 4362 753, 4258 219

Email: ncasc@mos.com.np

Fax: +977 - 1 - 4261 406

Website: www.ncasc.gov.np